



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : C07H 17/08, A61K 31/70, A61P 29/00		A2	(11) International Publication Number: WO 00/42055 (43) International Publication Date: 20 July 2000 (20.07.00)
(21) International Application Number: PCT/EP00/00163 (22) International Filing Date: 12 January 2000 (12.01.00) (30) Priority Data: MI99A000061 15 January 1999 (15.01.99) IT (71) Applicant (for all designated States except US): ZAMBON GROUP S.P.A. [IT/IT]; Via della Chimica, 9, I-36100 Vicenza (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): PELLACINI, Franco [IT/IT]; Via G. Balla 14, I-20151 Milano (IT). BOTTA, Daniela [IT/IT]; Via Valleggio 4, I-22100 Como (IT). ROMAGNANO, Stefano [IT/IT]; Via D. Scarlatti 33, I-20090 Buccinasco (IT). MORIGGI, Ermanno [IT/IT]; Largo Crivelli 5, I-21052 Busto Arsizio (IT). PRADELLA, Lorenzo [IT/IT]; Via A. Diaz 4, I-20063 Cernusco sul Naviglio (IT). (74) Agent: LONGONI, Alessandra; Zambon Group S.p.A., Corp. Patent & Trademark Dept., Via Lillo Del Duca, 10, I-20091 Bresso (IT).			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  Published Without international search report and to be republished upon receipt of that report.
(54) Title: MACROLIDES WITH ANTI-INFLAMMATORY ACTIVITY			
(57) Abstract  3'-Des-dimethylamino-9-oxyimino macrolides of formula (I) endowed with anti-inflammatory activity and their pharmaceutical use are described.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# "MACROLIDES WITH ANTI-INFLAMMATORY ACTIVITY"

\*\*\*\*\*

The present invention relates to macrolides with anti-inflammatory activity and, more particularly, it relates to des-dimethylamino macrolide derivatives with anti-inflammatory activity, their pharmaceutically acceptable salts and pharmaceutical compositions containing them as active ingredients.

It is known that many antibiotics, in particular the class of macrolides with 14 atoms derived from erythromycin, are endowed with anti-inflammatory properties in addition to the antibacterial activity [Clin. Immunother., (1996), 6, 454-464].

Erythromycin is a natural macrolide (The Merck Index, XII edition, n° 3720, page 625) that has had a very broad clinical use in the treatment of infections caused by Gram-positive bacteria, by some Gram-negative ones or by Mycoplasma.

Recently the interest of the scientific community has been focused on the anti-inflammatory and immunomodulatory component of erythromycin and derivatives [Journal of Antimicrobial Chemotherapy, (1998), 41, Suppl.B, 37-46].

Such activity is well documented by both clinical studies and *in vivo* and *in vitro* experiments.

For example, macrolides have proved to be effective in the therapy of inflammatory diseases such as panbronchiolitis [Thorax, (1997), 52, 915-918], bronchial asthma [Chest, (1991), 99, 670-673] and cystic fibrosis [The Lancet, (1998), 351, 420], or in animal models of inflammation such as, for example, the zymosan-induced peritonitis in mice [Journal of Antimicrobial Chemotherapy, (1992), 30, 339-348] and the neutrophil recruitment induced by endotoxin in rat trachea [Antimicrobial Agents and Chemotherapy, (1994), 38, 1641-1643] or in *in vitro* studies on immune system cells, such as neutrophils [The journal of Immunology, (1997), 159, 3395-4005] and T-lymphocytes [Life Sciences, (1992), 51, PL 231-236] or in the modulation of cytokines, such as interleukin 8 (IL-8) [Am. J. Respir. Crit. Care Med., (1997), 156, 266-271] or interleukin 5 (IL-5) (EP 0 775 489 and EP 0 771 564, Taisho Pharmaceutical Co., Ltd).

The peculiar therapeutic efficacy of macrolides on diseases in which the conventional anti-inflammatory drugs, such as for example corticosteroids, have demonstrated to be ineffective

-2-

[Thorax, (1997), 52, 915-918, already cited] justifies the high interest towards this new potential class of anti-inflammatories.


Nevertheless the strong antibacterial activity of the conventional macrolides does not allow an enlarged use in the chronic treatment of inflammatory processes not due to pathogens  
5 because of the rapid onset of resistant strains.

Therefore, it would be desirable to have new substances with a macrolide structure that show anti-inflammatory activities and, in the mean time, that are devoid of antibiotic properties.

For a greater clarity we show the formula of erythromycin in which the numbering adopted in the present patent application is indicated.

10

15



20


Some classes of erythromycin derivatives endowed with a high anti-inflammatory activity are described in the literature.

For example, in the already cited European patent applications in the name of Taisho derivatives of erythromycin modified at 3, 9, 11 and 12 position, as strong inhibitors of the synthesis of IL-5 are claimed.

25

N-alkyl derivatives of azithromycin, without cladinose and desosamine, of formula

-3-



10 wherein

R<sub>1</sub> is hydrogen, a lower alkyl or a lower alkanoyl; R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub>, the same or different, are hydrogen or a lower alkanoyl; are described as anti-inflammatories in EP 0 283 055 (Sour Pliva).


15 The use of erythromycin as anti-inflammatory that acts by reducing the release of interleukin I through the inhibition of the mammalian mdr-P glycoprotein is claimed WO 92/16226 in the name of Smith-Kline Beecham Corporation.

Among the macrolide derivatives described in the literature a few are 3-desdimethylamino-9-oxyimino derivatives. The limited interest towards this class of compounds is justified by the fact that the relevance of the dimethylamino group for the activity of ribosomal binding  
20 typical of macrolides is known [Tetrahedron Letters, (1994), 35, 3837-3840].

In US 3.928.387 (Hoffmann-La Roche Inc.) 3'-desdimethylamino-3',4'-dehydroerythromycin A oxime is described, as intermediate useful for the preparation of the antibiotic 1745A/X.

25 In EP 0 254 534 (Robinson, William S.) a very broad class of macrolides with antiviral activity is claimed. Among them the compound of formula


-4-



whose correct chemical name is 3'-desdimethylamino-3',4'-dehydroerythromycin A 9-O-methyloxime notwithstanding in the text of EP 0 254 534 is erroneously reported as desdimethylaminoerythromycin 9-O-methyloxime (page 10, line 46) is described.

15 Now we have found that by removing the dimethylamino group from the 3' position of desosamine of 9-oxyimino macrolides, compounds endowed with anti-inflammatory activity and essentially devoid of antibiotic properties are obtained.

Therefore, object of the present invention are compounds of formula




wherein

-5-

R is hydrogen or methyl;

R<sub>1</sub> and R<sub>2</sub> are both hydrogen or they together form a bond;

R<sub>3</sub> is hydrogen, a linear or branched C<sub>1</sub>-C<sub>5</sub> alkyl group, a benzyl group, optionally substituted by one or more substituents selected among nitro groups, hydroxy groups, carboxylic groups, amino groups, linear or branched C<sub>1</sub>-C<sub>5</sub> alkyl groups, C<sub>1</sub>-C<sub>4</sub> alkoxy carbonyl groups, aminocarbonyl groups or ciano groups, or a chain of formula



wherein

A is hydrogen or a phenyl group optionally substituted by one or two substituents selected among nitro groups, hydroxy groups, carboxylic groups, amino groups, linear or branched C<sub>1</sub>-C<sub>5</sub> alkyl groups, C<sub>1</sub>-C<sub>4</sub> alkoxy carbonyl groups, aminocarbonyl groups or ciano groups, or a 5 or 6 membered heterocycle, saturated or unsaturated, containing from 1 to 3 heteroatoms selected among nitrogen, oxygen and sulphur, optionally substituted by one or two substituents selected among C<sub>1</sub>-C<sub>5</sub> alkyl groups, phenyl groups, hydroxy groups, oxo (=O) groups, nitro groups, C<sub>1</sub>-C<sub>4</sub> alkoxy carbonyl groups, aminocarbonyl groups, mono or di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl groups, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl groups;

X and Y, the same or different, are O, S, SO, SO<sub>2</sub> or NR<sub>4</sub>, in which R<sub>4</sub> is hydrogen, a linear or branched C<sub>1</sub>-C<sub>5</sub> alkyl group, a C<sub>1</sub>-C<sub>5</sub> alkoxy carbonyl group, a benzyloxy carbonyl group;

r is an integer from 1 to 6;

m is an integer from 1 to 8;

n is an integer from 0 to 2;

and their pharmaceutically acceptable salts;

the compounds 3'-desdimethylamino-3',4'-dehydroerythromycin A oxime (R<sub>1</sub> and R<sub>2</sub>=bond; R=H; R<sub>3</sub>=H) and 3'-desdimethylamino-3',4'-dehydroerythromycin A 9-O-methyloxime (R<sub>1</sub> and R<sub>2</sub>=bond; R=H; R<sub>3</sub>=CH<sub>3</sub>) being excluded.

A further object of the present invention is the use of the compounds 3'-desdimethylamino-3',4'-dehydroerythromycin A oxime and 3'-desdimethylamino-3',4'-dehydroerythromycin A 9-O-methyloxime as anti-inflammatories.

The compounds of formula I are anti-inflammatory macrolides devoid of antibiotic activity

-6-


and therefore they are useful in the treatment of inflammatory diseases.

With the term linear or branched C<sub>1</sub>-C<sub>5</sub> alkyl groups a group selected among methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl and isopentyl is meant.

With the term 5 or 6 membered heterocycle, saturated or unsaturated, containing from 1 to 3  
 5 heteroatoms selected among nitrogen, oxygen and sulphur, heterocycles such as pyrrole, thiophene, furan, imidazole, pyrazole, thiazole, isothiazole, isoxazole, oxazole, pyridine, pyrazine, pyrimidine, pyridazine, triazole, thiadiazole and their partially or totally saturated forms are meant.

Preferred compounds of formula I are the compounds in which R, R<sub>1</sub> and R<sub>2</sub> are hydrogen.

10 Within this class the compounds in which R<sub>3</sub> is a chain of formula




wherein

X, Y, A, *r*, *m* and *n* have the already reported meanings are particularly preferred.

Even more preferred compounds are the compounds in which R<sub>3</sub> is a chain of formula

15



wherein

*r* is 2, *m* is 2 or 6, *n* is 1, Y is NR<sub>4</sub>, X is O or NR<sub>4</sub>, R<sub>4</sub> is hydrogen and A is phenyl or thiazolyl.

Examples of pharmaceutically acceptable salts of the compounds of formula (I) are the salts  
 20 with organic or inorganic acids such as hydrochloric, hydrobromic, hydriodic, nitric, sulphuric, phosphoric, acetic, tartaric, citric, benzoic, succinic and glutaric acid.

The compounds of formula I, object of the present invention, are prepared by following a synthetic scheme which comprises (a) the removal of the dimethylamino group at 3' position and (b) the optional functionalisation of the oxime.


25 The removal of the dimethylamino group is performed by oxidation, pyrolysis and optional reduction, according to known methods. It is evident to the man skilled in the art that, in order to avoid interferences with the functional groups optionally present at R<sub>3</sub> substituent, the removal of the dimethylamino group will be preferably accomplished starting from intermediates of formula



-7-

5

10



(II)


in which

R has the already reported meanings and R<sub>3</sub>' is hydrogen or a linear or branched C<sub>1</sub>-C<sub>5</sub> alkyl group.

By oxidation the corresponding N-oxides of formula

15

20



(III)


25 in which

R and R<sub>3</sub>' have the already reported meanings are obtained;

that by pyrolysis, optionally followed by reduction, respectively give the compounds of formula


-8-

5

(I-A;  $R_1$  and  $R_2$ =bond)

and

10

(I-B;  $R_1=R_2=H$ )

15

20 object of the present invention (I- $R_3$ =hydrogen or  $C_1$ - $C_5$  alkyl).

According to common techniques, the compounds of formula I in which  $R_3$  is different from hydrogen can be prepared from the compounds of formula I-A and I-B in which  $R_3'$  is hydrogen by functionalisation of the oxime.

Generally the functionalisation is carried out by reaction with a compound of formula

25


 $R_3''$ -W

(IV)

in which  $R_3''$  has all the meanings of  $R_3$  apart from hydrogen and W is a leaving group, preferably a chlorine or bromine atom or a mesyl group.

An alternative synthetic route particularly suited for the preparation of the compounds of formula I in which  $R_3$  is a chain of formula


-9-



in which

X, Y, A,  $r$ ,  $m$  and  $n$  have the already reported meanings;


- 5 comprises the reaction of a compound of formula I in which  $R_3$  is hydrogen with an intermediate of formula



(V)

in which


- 10 W, X, Y,  $m$  and  $n$  have the already reported meanings and Z represents a protecting group;  
to give the intermediate of formula



(VI)

in which R,  $R_1$ ,  $R_2$ , X, Y, Z,  $r$  and  $m$  have the already reported meanings;

that after removal of the Z protecting group, is reacted with a derivative of formula



(VII)

in which A, W and  $n$  have the already reported meanings;

to give the compounds of formula I.

The compounds of formula I in which Y is  $\text{NR}_4$  can be prepared according to the above reported synthetic route also by using an aldehyde of formula

-10-

A-CHO

(VIII)

in which A has the already reported meanings;

in place of the intermediate of formula VII, subject to removal of the Z protecting group from the intermediate of formula VI.

- 5 Moreover, the compounds of formula I in which  $R_1=R_2=H$  can be prepared by reduction of the corresponding compounds of formula I in which  $R_1$  and  $R_2$  form a bond.

The compounds of formula I, object of the present invention, are endowed with an anti-inflammatory activity and are devoid of antibiotic activity.

- 10 The pharmacological activity of the compounds of formula I has been evaluated by *in vitro* and *in vivo* tests in comparison with known macrolides, such as erythromycin, clarithromycin and roxithromycin, endowed with both anti-inflammatory activity and antibiotic activity.

- 15 The anti-inflammatory activity has been evaluated *in vitro* as inhibition of IL-8 release and of superoxide anion release (example 11) and *in vivo* as inhibition of LPS-induced neutrophilia after repeated administrations (example 12).

In all the experiments the compounds object of the present invention have resulted very active as anti-inflammatories and the anti-inflammatory activity has been equal to or greater than the one of the reference compounds.

- 20 For a therapeutical application the compound of formula I can be used in a pharmaceutical form suitable to oral or parenteral administration.

Therefore, object of the present invention are pharmaceutical compositions containing a therapeutically active amount of a compound of formula I or of a salt thereof in admixture with a pharmaceutically acceptable carrier.

With the aim to better illustrate the present invention the following examples are now given.

- 25 Example 1

Preparation of [6-(2-hydroxy-ethylamino)-hexyl]-carbamic acid benzyl ester

To a solution of (6-hydroxy-hexyl)-carbamic acid benzyl ester (25 g; 99.47 mmol), prepared as described in WO 96/18633, in  $CH_2Cl_2$  (350 ml), cooled with ice at about 10°C, at first a solution of KBr (1.18 g; 9.94 mmol) in water (20 ml) and TEMPO (0.155 g; 0.994

-11-

mmoles) and then, dropwise in about 15-20 minutes keeping the temperature at 10-12°C, a solution prepared with NaHCO<sub>3</sub> (7.5 g; 89.28 mmoles) and NaClO (4.5 % water solution; 197 ml; 125 mmoles) were added.

15 minutes after the end of the dripping, the phases were separated and the aqueous phase  
5 was extracted once with CH<sub>2</sub>Cl<sub>2</sub> (100 ml). The collected organic extracts were washed twice with brine (20% NaCl) and dried on sodium sulphate.

3 Å molecular sieves (30 g) and then, by fast dropwise and by cooling with water and ice, a solution of 2-aminoethanol (35.9 ml; 0.597 moles) in ethanol (600 ml) were added to the obtained solution (about 800 ml).

10 Ended the dripping the mixture was kept under stirring at room temperature for 2 hours and then filtered.

NaBH<sub>4</sub> (4.54 g; 120 mmoles) was portionwise added to the obtained solution, cooled with water and ice, under stirring in a nitrogen atmosphere.

At the end of the addition, the reaction mixture was kept under stirring at room temperature  
15 for 2 hours and then the solvent was evaporated.

The residue was collected with water and ethyl acetate and the phases were separated, again extracting twice the aqueous phase with ethyl acetate.

The collected organic extracts were washed with brine (20% NaCl), dried on sodium sulphate and concentrated up to obtained an oily residue that was inclined to solidify.

20 The residue was triturated with hexane, filtered and washed with a mixture of hexane and ethyl ether, furnishing [6-(2-hydroxy-ethylamino)-hexyl]-carbamic acid benzyl ester (26.22 g; yield 89%) as a white solid.

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.33-7.25 (m, 5H, Ar); 5.05 (s, 2H, COOCH<sub>2</sub>); 4.96 (broad-t, 1H, NH); 3.63-3.58 (m, 2H, \*CH<sub>2</sub>-OH); 3.19-3.09 (m, 2H, CH<sub>2</sub>NCO); 2.72-2.67  
25 (m, N-\*CH<sub>2</sub>-CH<sub>2</sub>O); 2.59-2.52 (m, 4H, OH and CH<sub>3</sub>); 1.53-1.23 (m, 8H, 4CH<sub>2</sub>).

### Example 2

#### Preparation of 6-(benzyloxycarbonylamino-hexyl)-(2-hydroxy-ethyl)-carbamic acid benzyl ester

A solution of benzylchloroformate benzyl ester (50% in toluene; 42.5 ml; 0.128 moles) in

-12-

ethyl acetate (85.5 ml) and 1N NaOH (128 ml; 0.128 moles) were simultaneously dropwise added to a solution of [6-(2-hydroxy-ethylamino)-hexyl]-carbamic acid benzyl ester (31.5 g; 0.107 moles), prepared as described in example 1, in a mixture of water (87 ml), 1N NaOH (17 ml) and ethyl acetate (180 ml), cooled at 0-5°C, controlling temperature and pH (about 5 8).

At the end of the dripping, the reaction mixture was kept under stirring for 30 minutes at 0-5°C, then the cooling was removed and further 1N NaOH (15 ml) was added, in order to bring again the pH at 8, then leaving under stirring at room temperature overnight.

The phases were separated and the aqueous phase was extracted once again with ethyl 10 acetate. The collected organic extracts were washed with brine, dried on sodium sulphate and concentrated under vacuum up to obtain an oily residue.

By chromatographical purification (eluant ethyl acetate: petrolatum from 60:40 to 70:30) 6-(benzyloxycarbonylamino-hexyl)-(2-hydroxy-ethyl)-carbamic acid benzyl ester (42.5 g; yield 92%) as an oil was obtained.

15 <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.39-7.25 (m, 10H, Ar); 5.10 and 5.07 (2s, 4H, 2COOCH<sub>2</sub>); 3.71 (broad signal, 2H, \*CH<sub>2</sub>-OH); 3.43-3.01 (m, 4H, 2CH<sub>2</sub>NCO); 1.57-1.19 (m, 8H, 4CH<sub>2</sub>).

By working in a similar way the following compounds were obtained:

(2-benzyloxycarbonylamino-ethyl)-(2-hydroxy-ethyl)-carbamic acid benzyl ester

20 starting from 2-(2-aminoethylamino)-ethanol.

(yield 32%)

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.33-7.28 (m, 10H, 2Ph); 5.06 and 5.04 (2s, 4H, 2CH<sub>2</sub>-Ph); 3.73-3.34 (broad m, 8H, 4CH<sub>2</sub>).

[2-(benzyl-benzyloxycarbonyl-amino)-ethyl]-(2-hydroxy-ethyl)-carbamic acid benzyl ester

25 starting from 2-[2-(benzylamino)-ethylamino]-ethanol prepared as described in WO 96/18633.

(yield 50%)

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): (very broad signals) 7.42-7.13 (m, 15H, 3Ar); 5.12 and 5.09 (2s, 4H, COOCH<sub>2</sub>\*); 4.55 (s, 2H, N-\*CH<sub>2</sub>-Ph).

-13-

[2-(2-hydroxy-ethoxy)-ethyl]-carbamic acid benzyl ester

starting from 2-(2-aminoethoxy)-ethanol

(yield 85%)

- <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.36-7.28 (m, 5H, Ar); 5.18 (broad signal, 1H, NH);  
 5.08 (s, 2H, Ph-\*CHO); 3.74-3.34 (m, 8H, \*CH<sub>2</sub>-\*CH<sub>2</sub>-O-\*CH<sub>2</sub>-\*CH<sub>2</sub>-OH); 2.13 (broad-t, 1H, OH).

## Example 3

Preparation of methansulphonic acid 2-[benzyloxycarbonyl-(6-benzyloxycarbonylamino-hexyl)-amino]-ethyl ester

- 10 Triethylamine (8.95 ml; 64.31 mmol) was added to a solution of 6-(benzyloxycarbonylamino-hexyl)-(2-hydroxy-ethyl)-carbamic acid benzyl ester (13.78 g; 32.15 mmol), prepared as described in example 2, in CH<sub>2</sub>Cl<sub>2</sub> (140 ml). The mixture was cooled at 0-5°C and then was dropwise added to a solution of methansulphonylchloride (3.36 ml; 43.41 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 ml).
- 15 At the end of the addition, the mixture was kept under stirring at room temperature for 60 minutes, then washed with 5% aqueous citric acid, with brine (20% NaCl), with 5% aqueous NaHCO<sub>3</sub> and finally with brine again. After drying on sodium sulphate and evaporation under vacuum methansulphonic acid 2-[benzyloxycarbonyl-(6-benzyloxycarbonylamino-hexyl)-amino]-ethyl ester (16.37 g; yield 100%) as a brown oil was obtained.
- 20 <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.35-7.27 (m, 10H, Ar); 5.11 and 5.07 (2s, 4H, 2COOCH<sub>2</sub>); 4.36-4.19 (m, 2H, CH<sub>2</sub>OSO<sub>2</sub>); 3.57-3.51 (m, 2H, SO-CH<sub>2</sub>-\*CH<sub>2</sub>N); 3.32-3.07 (m, 4H, 2CH<sub>2</sub>N); 2.91 and 2.85 (2s-conformers, 3H, CH<sub>3</sub>); 1.50-1.20 (m, 8H, 4CH<sub>2</sub>).

By operating in a similar way the following compounds were obtained:

methansulphonic acid 2-[2-(benzyloxycarbonyl-amino)-ethoxy]-ethyl ester

- 25 starting from [2-(2-hydroxy-ethoxy)-ethyl]-carbamic acid benzyl ester, prepared as described in example 2.
- (yield 98%)

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.36-7.28 (m, 5H, Ar); 5.16 (broad signal, 1H, NH); 5.08 (s, 2H, COOCH<sub>2</sub>); 4.34-4.30 (m, 2H, SO<sub>3</sub>CH<sub>2</sub>); 3.72-3.67 (m, 2H, SO<sub>3</sub>-CH<sub>2</sub>-\*CH<sub>2</sub>);

-14-

3.60-3.34 (m, 4H, N-CH<sub>2</sub>-CH<sub>2</sub>); 2.98 (s, 3H, SO<sub>3</sub>CH<sub>3</sub>).

methansulphonic acid 2-[[2-(benzyl-benzyloxycarbonyl-amino)-ethyl]-benzyloxycarbonyl-amino]-ethyl ester

starting from [2-(benzyl-benzyloxycarbonyl-amino)-ethyl]-(2-hydroxy-ethyl)-carbamic acid  
5 benzyl ester, prepared as described in example 2.

(yield 72%)

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.40-7.00 (m, 15H, 3Ar); 5.13-5.01 (broad signal, 4H, 2COOCH<sub>2</sub>); 4.51-3.30 (broad-m, 10H, 4CH<sub>2</sub>N and CH<sub>2</sub>SO<sub>3</sub>); 2.92-2.76 (broad signal, 3H, CH<sub>3</sub>).

10 methansulphonic acid 2-[benzyloxycarbonyl-(2-benzyloxycarbonylamino-ethyl)-amino]-ethyl ester

starting from (2-benzyloxycarbonylamino-ethyl)-(2-hydroxy-ethyl)-carbamic acid benzyl ester, prepared as described in example 2.

(yield 100%)

15 <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.35-7.27 (m, 10H, 2Ar); 5.10 and 5.04 (2s, 4H, 2COOCH<sub>2</sub>); 4.41-4.15 (m, 2H, CH<sub>2</sub>-MeSO<sub>2</sub>); 3.63-3.23 (m, 6H, N-CH<sub>2</sub>-CH<sub>2</sub>-N-CH<sub>2</sub>); 2.90 (s-broad, 3H, MeSO<sub>2</sub>).

#### Example 4

##### Preparation of erythromycin A oxime N-oxide

20 A solution of H<sub>2</sub>O<sub>2</sub> (72.00 g; titre 34% w/v; 0.72 moles) in water (780 ml) was dropwise added in 1 hour to a solution of erythromycin A oxime (35.00 g; 0.0467 moles) in methanol (1400 ml) under mechanical stirring, keeping the temperature at 20-25°C. At the end of the addition the reaction mixture was kept under stirring and at room temperature for 24 hours.  
After having added more H<sub>2</sub>O<sub>2</sub> (8 ml), the mixture was kept under stirring for further 6  
25 hours.

Methanol was evaporated under vacuum at a temperature of about 40°C maintaining the volume of water constant (about 700 ml).

After filtering, washing with water and drying erythromycin A oxime N-oxide (36.3 g; yield 99%) was obtained as a white crystal solid.



-15-

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>) δ (ppm): 10.71-10.19 (broad signal, 2H, shifting H); 5.14-5.08 (m, 1H, H-13); 4.72 (d, 1H, J<sub>HH</sub>=4.4 Hz, H-1''); 4.45 (d, 1H, J<sub>HH</sub>=7.0 Hz, H-1').

#### Example 5

##### Preparation of 3'-de(dimethylamino)-3',4'-dehydro-erythromycin A oxime (Compound 1)

5 A solution of erythromycin A oxime N-oxide (30.00g; 38.3 mmoles), prepared as described in example 4, in dimethylformamide (235 ml) was warmed at 150°C in a pre-warmed oil bath (175-180°C) and left at such temperature under mechanical stirring for 15-20 minutes.

After cooling and evaporation of dimethylformamide, the oily residue was collected with demineralized water (500 ml), warmed and cooled. The filtered solid was triturated and dried  
10 under vacuum at 40-45°C furnishing a crude (25.5 g).

The crude was at first crystallized from acetonitrile (110 ml), filtered, washed with water and dried under vacuum at 50°C obtaining a crystal product (20 g) which was crystallized again from methanol/water=65/35 (400 ml), filtered and dried under vacuum at 40-50°C.

Compound 1 as a crystal product was so obtained (10.3 g; yield 38.2%).

15 More product (3.7 g) was then recovered from the crystallization liquor with an overall yield of 51.8%.

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 5.67-5.55 (m, 2H, \*CH=\*CH); 4.44 (d, 1H, J<sub>HH</sub>=7.0 Hz, H-1'); 4.33-4.22 (m, 1H, H-5'); 4.13-4.04 (m, 1H, H-2'); 3.84-3.73 (m, 1H, H-8); 3.69 (s, 1H, H-11).

20 <sup>13</sup>C-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 171.11 (s, C-9); 132.2 and 126.1 (2s, C-3' and C-4').

#### Example 6

##### Preparation of 3'-de(dimethylamino)-erythromycin A oxime (Compound 2)

Platinum oxide (0.615 g) was added at room temperature to a solution of compound 1 (20.00 g; 28.4 mmoles), prepared as described in example 5, in ethanol (850 ml) (the complete  
25 dissolution was obtained after slight heating).

The mixture was hydrogenated in a Parr apparatus (1.36 atm) and the absorption was immediate.

After filtration of the catalyst and evaporation of the solvent under vacuum, the white crystal residue was triturated with petrolatum, filtered and dried under vacuum at 50°C furnishing

-16-

compound **2** (19.8 g; yield 99%).

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 5.05-4.98 (m, 1H, H-13); 4.94 (d, 1H, J<sub>HH</sub>=4.4 Hz, H-1''); 4.23 (d, 1H, J<sub>HH</sub>=7.4 Hz, H-1'); 3.44-3.30 (m, 1H, H-2'); 2.07-1.21 (m, 2H, H-3'); 1.65-1.45 (m, 2H, H-4').

5 <sup>13</sup>C-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 171.2 (s, C-9); 104.7 (s, C-1'); 31.8 (s, C-3'); 29.5 (s, C-4').

#### Example 7

Preparation of 3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[benzyloxy-carbonyl-(6-benzyloxy-carbonylamino-hexyl)-amino]-ethyl]-oxime] (Compound 3)

10 Compound **2** (12.51 g; 17.73 mmol), prepared as described in example 6, was added to a 95% solution of potassium tert-butyrate (2.45 g; 19.48 mmol) in anhydrous THF (120 ml), under stirring and in a nitrogen atmosphere, maintaining the temperature at about 25°C. The reaction mixture was kept under stirring for 30 minutes at room temperature and then 18-crown-6 ether (4.69 g; 17.73 mmol) and a solution of methansulphonic acid 2-  
15 [benzyloxycarbonyl-(6-benzyloxycarbonylamino-hexyl)-amino]-ethyl ester (8.98 g; 17.73 mmol), prepared as described in example 3, in anhydrous THF (60 ml), were added, leaving under stirring at room temperature overnight.

After evaporation of the solvent under vacuum, the residue was taken up with a mixture of ethyl acetate and brine (20% NaCl) and the phases were separated. The aqueous phase was  
20 again extracted with ethyl acetate. The collected and dried organic extracts were concentrated under vacuum obtaining a crude (23.4 g).

By chromatographycal purification (eluant CH<sub>2</sub>Cl<sub>2</sub>:CH<sub>3</sub>OH=97:3) still slightly impure compound **3** was obtained (15.42 g), which was used without further purifications.

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.35-7.28 (m, 10H, Ar); 5.14-5.06 (m, 1H, H-13);  
25 5.11 and 5.07 (2s, 4H, 2COOCH<sub>2</sub>); 4.84 (d, 1H, J<sub>HH</sub>=4.4 Hz, H-1''); 4.28 (d, 1H, J<sub>HH</sub>=7.4 Hz, H-1').

By working in a similar way the following compounds were obtained:

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[2-(benzyloxycarbonylamino)-ethoxy]-ethyl]-oxime] (Compound 4) from methansulphonic acid 2-[2-(benzyloxycarbonyl-amino)-

-17-

ethoxy]-ethyl ester.

(yield 62%)

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.34-7.27 (m, 5H, Ar); 6.09 (broad signal, 1H, NH); 5.13-5.05 (m, 1H, H-13); 5.06 (s, 2H, COOCH<sub>2</sub>); 4.79 (d, 1H, J<sub>HH</sub>=4.4 Hz, H-1''); 4.26 (d, 1H, J<sub>HH</sub>=7.4 Hz, H-1').

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-methoxy-ethoxy]-methyl]-oxime]

(Compound 5) from methoxy-ethoxy-methyl chloride

(yield 32.5%)

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 5.21-5.12 (m, 2H, O-CH<sub>2</sub>-O); 5.12-5.05 (m, 1H, H-13); 4.85 (d, 1H, J<sub>HH</sub>=5.4 Hz, H-1''); 4.39 (d, 1H, J<sub>HH</sub>=7.5 Hz, H-1'); 3.40 (s, 3H, CH<sub>2</sub>-O-CH<sub>3</sub>); 3.27 (s, 3H, H-3').

<sup>13</sup>C-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 172.5 (s, C-9); 97.44 (s, NO-C-O); 32.12 (s, C-3'); 29.84 (s, C-4').

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[[2-(benzyl-benzyloxycarbonyl-amino)-ethyl]-benzyloxycarbonyl-amino]-ethyl]-oxime] (Compound 12) from methansulphonic acid 2-[[2-(benzyl-benzyloxycarbonyl-amino)-ethyl]-benzyloxycarbonyl-amino]-ethyl ester

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.40-7.16 (broad m., 15H, 3Ph); 5.17-5.00 (m, 5H, 2\*CH<sub>2</sub>-Ph and H-13).

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[[2-(benzyloxycarbonyl-amino)-ethyl]-benzyloxycarbonyl-amino]-ethyl]-oxime] (Compound 13) from methansulphonic acid 2-[benzyloxycarbonyl-(2-benzyloxycarbonylamino-ethyl)-amino]-ethyl ester

(yield 42%)

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.38-7.25 (m, 10H, 2Ph); 5.11 and 5.05 (2s, 4H, 2COOCH<sub>2</sub>); 5.14-5.00 (m, 1H, H-13); 4.89-4.79 (broad-m, 1H, H-1''); 2.26 (d, 1H, J<sub>HH</sub>=7.4 Hz, H-1').

#### Example 8

Preparation of 3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[(6-amino-hexyl)-amino]-ethyl]-oxime] (Compound 6)

10% Pd/C (1.6 g) was added to a solution of compound 3 (15.42 g; 13.8 mmoles), obtained

-18-

as described in example 7, in ethanol (160 ml).

The mixture was hydrogenated in a Parr apparatus (1.02 atm). After 2 hours the catalyst was filtered and the solvent was evaporated.

The residue was purified by flash chromatography. (eluant  $\text{CH}_2\text{Cl}_2:\text{CH}_3\text{OH}:\text{NH}_3=85:15:1.5$   
5 then 80:20:2) furnishing compound 6 (8.48 g) as an amorphous white solid.

$^1\text{H}$ -NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm): 5.07-5.00 (m, 1H, H-13); 4.79 (d, 1H,  $J_{\text{HH}}=4.4$  Hz, H-1''); 4.21 (d, 1H,  $J_{\text{HH}}=7.4$  Hz, H-1').

$^{13}\text{C}$ -NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm): 171.8 (s, C-9); 71.6 (s, =N-O-C); 49.43 and 49.0 (2s, =N-O-C-\*C-N-\*C); 41.1 (s, C-NH<sub>2</sub>).

10 By working in a similar way the following compounds were obtained:

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-(2-amino-ethoxy)-ethyl]-oxime]

(Compound 7) starting from compound 4

(yield 85%)

$^1\text{H}$ -NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm): 5.12-5.05 (m, 1H, H-13); 4.83 (d, 1H,  $J_{\text{HH}}=4.4$  Hz, H-1''); 4.26 (d, 1H,  $J_{\text{HH}}=7.5$  Hz, H-1').  
15

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[(2-benzylamino-ethyl)-amino]-ethyl]-oxime] (Compound 14) starting from compound 12

(yield 48%)

$^1\text{H}$ -NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm): 7.32-7.17 (m, 5H, Ph); 5.08-5.00 (m, 1H, H-13); 4.74 (d, 1H,  $J_{\text{HH}}=4.6$  Hz, H-1''); 4.22 (d, 1H,  $J_{\text{HH}}=7.4$  Hz, H-1'); AB system:  $V_a=3.80$ ,  $V_b=3.76$ ,  $J_{ab}=13.7$  Hz, \*CH<sub>2</sub>Ph.  
20

$^{13}\text{C}$ -NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm): 171.3 (s, C-9); 105.0 (s, C-1'); 32.2 (s, C-3'); 29.8 (s, C-4').

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[(2-amino-ethyl)-amino]-ethyl]-oxime]

25 (Compound 15) starting from compound 13

(yield 70%)

$^1\text{H}$ -NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm): 5.08-5.00 (m, 1H, H-13); 4.76 (d, 1H,  $J_{\text{HH}}=4.6$  Hz, H-1''); 4.23 (d, 1H,  $J_{\text{HH}}=7.4$  Hz, H-1').

Example 9

Preparation of 3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[6-[(thiazol-2-ylmethyl)-amino]-hexyl-amino]-ethyl]-oxime] (Compound 8)

A suspension of compound 6 (3 g; 3.53 mmol), prepared as described in example 8, 97% 2-thiazolecarbaldehyde (0.412 g; 3.53 mmol) and molecular sieves (3 Å, 6.75 g) in ethanol  
5 (60 ml) was left under stirring for 3 hours.

After filtering the molecular sieves on celite, 10% Pd/C (0.3 g) was added and the mixture was hydrogenated in a Parr hydrogenator (1.02 atm). After 20 hours the catalyst was filtered and the solvent evaporated.

By chromatographical purification of the residue (eluant  
10 CHCl<sub>3</sub>:petrolatum:triethylamine=90:10:10) compound 8 was obtained (1.66 g; yield 49.8%) as a amorphous solid.

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.66 (d, 1H, J<sub>HH</sub>=3.2 Hz, CHN); 7.22 (d, 1H, CHS); 5.08-5.02 (m, 1H, H-13); 4.78 (d, 1H, J<sub>HH</sub>=4.4 Hz, H-1''); 4.21 (d, 1H, J<sub>HH</sub>=7.4 Hz, H-1'); 4.07 (s, 2H, \*CH<sub>2</sub>-thiaz.).

15 <sup>13</sup>C-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 172.0 (s, S-C=N); 171.7 (s, C-8); 142.4 (s, CHN); 118.7 (s, CHS); 104.9 (s, C-1'); 96.3 (s, C-1''); 71.7 (s, N-O-C).

By working in a similar way the following compounds were obtained:

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[6-(benzylamino)-hexylamino]-ethyl]-oxime] (Compound 9) from compound 6 and benzaldehyde.

20 <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.27-7.17 (m, 5H, Ar); 5.07-5.01 (m, 1H, H-13); 4.75 (d, 1H, J<sub>HH</sub>=4.4 Hz, H-1''); 4.19 (d, 1H, J<sub>HH</sub>=7.4 Hz, H-1'); 3.73 (s, 2H, \*CH<sub>2</sub>-Ph).

<sup>13</sup>C-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 171.8 (s, C-9); 104.9 (s, C-1'); 96.3 (s, C-1''); 71.8 (s, =N-O-C); 53.8 (s, N-\*C-Ph); 49.7, 49.2 and 49.1 (3s, 3N-C).

25 3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[2-[(thiazol-2-ylmethyl)-amino]-ethoxy]-ethyl]-oxime] (Compound 10) from compound 7 and 2-thiazolecarbaldehyde.

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.64 (d, 1H, J<sub>HH</sub>=3.2 Hz, CHN); 7.19 (d, 1H, CHS); 5.10-5.02 (m, 1H, H-13); 4.78 (d, 1H, J<sub>HH</sub>=4.4 Hz, H-1''); 4.21 (d, 1H, J<sub>HH</sub>=7.4 Hz, H-1'); 4.13 (s, 2H, \*CH<sub>2</sub>-thiaz.).

<sup>13</sup>C-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 172.7 (s, SC=N); 171.5 (s, C-9); 142.4 (s, CHN');

-20-

118.6 (s, CHS); 104.7 (s, C-1'); 96.4 (s, C-1''); 50.7 (s, N-\*C-thiaz.).

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[2-(benzylaminol)-ethoxy]-ethyl]-oxime]

(Compound 11) from compound 7 and benzaldehyde.

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.33-7.10 (m, 5H, Ar); 5.12-5.04 (m, 1H, H-13); 4.78  
5 (d, 1H, J<sub>HH</sub>=4.4 Hz, H-1''); 4.72 (d, 1H, J<sub>HH</sub>=7.4 Hz, H-1'); 3.80 (s, 2H, NCH<sub>2</sub>).

<sup>13</sup>C-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 171.5 (s, C-9); 104.8 (s, C-1'); 96.5 (s, C-1''); 69.4, 70.8 and 72.4 (3s, 3OCH<sub>2</sub>); 53.6 (s, \*C-Ph); 48.2 (s, O-C-\*C-N).

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[2-[(thiazol-2-ylmethyl)-amino]-ethyl]-amino]-ethyl]-oxime] (Compound 16) from compound 15 and 2-thiazolecarbaldehyde.

10 <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.62 (d, 1H, J<sub>HH</sub>=3.0 Hz, N-\*CH=CH); 7.18 (d, 1H, S-\*CH=CH); 4.18 (d, 1H, J<sub>HH</sub>=7.4 Hz, H1').

<sup>13</sup>C-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 172.6 (s, SC=N); 171.3 (s, C-9); 142.4 (s, CHN); 118.6 (s, CHS); 104.9 (s, C-1'); 96.4 (s, C-1''); 32.17 (s, C-3'); 29.8 (s, C-4').

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[6-[(2-phenyl-1H-imidazol-4-ylmethyl)-amino]-hexyl-amino]-ethyl]-oxime] (Compound 17) from compound 6 and 2-phenyl-1H-imidazole-4-carbaldehyde.

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.86-7.21 (m, 5H, Ar); 6.91 (s, 1H, CH-Imid.); 5.11-5.02 (m, 1H, H13); 4.76 (d, 1H, J<sub>HH</sub>=4.2 Hz, H1''); 4.21 (d, 1H, J<sub>HH</sub>=7.4 Hz, H1'); 3.76 (s, 2H, \*CH<sub>2</sub>-Imid.); 3.23 (s, 3H, OMe).

20 <sup>13</sup>C-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 171.7 (s, C-9); 104.8 (s, C-1'); 96.3 (s, C-1''); 32.1 (s, C-3'); 29.9 (s, C-4').

3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[6-[(1-methyl-2-phenyl-1H-imidazol-4-ylmethyl)-amino]-hexyl-amino]-ethyl]-oxime] (Compound 18) from compound 6 and 1-methyl-2-phenyl-1H-imidazole-4-carbaldehyde.

25 <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 7.57-7.31 (m, 5H, Ar); 6.87 (s, 1H, CH-Imid.); 5.09-5.00 (m, 1H, H13); 4.76 (d, 1H, J<sub>HH</sub>=4.2 Hz, H1''); 4.20 (d, 1H, J<sub>HH</sub>=7.4 Hz, H1'); 3.71 (s, 2H, \*CH<sub>2</sub>-Imid.); 3.62 (s, 3H, NMe); 3.21 (s, 3H, OMe).

<sup>13</sup>C-NMR (200 MHz, CDCl<sub>3</sub>) δ (ppm): 171.8 (s, C-9); 104.9 (s, C-1'); 96.3 (s, C-1''); 32.1 (s, C-3'); 29.7 (s, C-4').

-21-

## Example 10

Preparation of 3'-de(dimethylamino)-erythromycin A (E)-9-[O-[2-[2-[(thiazol-2-ylmethyl)-(methyl)-amino]-ethoxy]-ethyl]-oxime] (Compound 19)

Compound 10 (0.23 g; 0.258 mmoles), formaldehyde (37% w/v; 42  $\mu$ l; 0.516 mmoles), 10%  
5 Pd on charcoal (25 mg) and a 4:1 mixture of ethanol and water (10 ml) were charged in a Parr apparatus at 1.02 atm. After 2 and 3 hours further formaldehyde was added (42  $\mu$ l + 21  $\mu$ l) At the end of the reaction (overall 6 hours) the catalyst was filtered off and the solvent was evaporated. The resultant residue was purified by flash chromatography (eluant  $\text{CH}_2\text{Cl}_2:\text{CH}_3\text{OH}=95:5$ ) furnishing compound 19 as an amorphous solid.

10  $^1\text{H-NMR}$  (200 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm): 7.64 (d, 1H,  $\text{J}_{\text{HH}}=3.6$  Hz, N-\*CH=CH); 7.21 (d, 1H, S-\*CH=CH); 5.10-5.00 (m, 1H, H13); 4.80 (d, 1H,  $\text{J}_{\text{HH}}=4.6$  Hz, H1"); 4.23 (d, 1H,  $\text{J}_{\text{HH}}=7.4$  Hz, H1'); 3.94 (s, 2H, \*CH<sub>2</sub>-Thiaz.).

$^{13}\text{C-NMR}$  (200 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm): 172.0 (s, SC=N); 171.8 (s, C-9); 142.2 (s, CHN); 119.3 (s, CHS); 104.9 (s, C-1'); 96.4 (s, C-1"); 32.2 (s, C-3'); 29.9 (s, C-4').

15

## Example 11

In vitro pharmacological activity

A) Release of interleukin 8 (IL-8)

A human endothelial immortalized cell line (ECV304) was obtained from ATTC (Rockville, Md) and grown in Medium 199, mod. Earle's salts (GIBCO, Life  
20 Technologies, Grand Island, N.Y.), supplemented with 20% Foetal Calf Serum (GIBCO), 100 U/ml of penicillin and 100  $\mu$ g/ml of streptomycin (SIGMA, St. Louis, MO) in wet atmosphere with 5%  $\text{CO}_2$  at 37°C.

The cells were cultivated on 96 well plates up to obtain a confluent monolayer.

The compounds to be evaluated were dissolved in DMSO at  $10^{-2}\text{M}$  and diluted with  
25 culture medium.

The compounds were pre-incubated with the cells 1 hour before the challenge.

The release of IL-8 was induced by adding 0.66  $\mu$ g/ml of lipopolysaccharide B (E. coli 055:B5, Difco, Detroit, Mi) in a final volume of 200  $\mu$ l.

After one night, the supernatant was collected for the IL-8 test.



-22-

The specific immunoreactivity for IL-8 in the culture supernatant was measured with ELISA kit (Amersham, UK).

Results were expressed as the highest obtainable inhibition (efficacy) and, whether possible, as the concentration at which 50% of such effect ( $IC_{50}$ ) is obtained.

5    B)    Release of superoxide anions

10    Neutrophils were separated from venous blood of healthy volunteers by centrifugation on Ficoll-Hypaque, followed by sedimentation on 6% dextrane and osmotic lysis of erythrocytes. Neutrophils were then washed and resuspended in a medium made of RPMI-1640 supplemented with 5% foetal calf serum and 1.34 mmoles/l of disodium dihydrate EDTA. The cells were maintained for 24 hours at 4°C and before the test the suspension was centrifugated and resuspended in HBSS (Hanks' balanced salt solution). The vitality and the purity of the neutrophil preparation was verified by dyeing with Trypan blue and Turk blue.

15    The superoxide anions were measured by using Lucigenin (bis-N-methylacridinium nitrate)-enhanced chemiluminescence technique.

Neutrophils ( $2 \times 10^6$  cells/ml) were pre-incubated for 30 minutes at 37°C in 900  $\mu$ l of HBSS with and without the tested compound (reading system).

20    The production of superoxide anions was measured with a Lumac/3M bio-counter after addition of 100  $\mu$ l of a solution in HBSS of Lucigenin (2 mmoles/l) and N-formyl-L-methionyl-L-leucyl-L-phenylalanine (FLMP) as stimulating agent at the concentration of  $1 \times 10^{-5}$  M to the reading system.

25    FLMP was dissolved in DMSO ( $1 \times 10^{-2}$  M) and further diluted in HBSS. The compound to be tested was dissolved in DMSO at the concentration of  $10^{-2}$  M. Lower concentrations, in DMSO, were prepared from that solution, and tested. The DMSO amount present in the reading system was lower than 1%.

The luminosity values obtained at the peak, for each tested concentration of the compound, were transformed in inhibition percentage in comparison with the reference.

The concentration able to inhibit the production of superoxide anion at 50% ( $IC_{50}$ ) was



-23-

calculated from the obtained "dose-response" curve.

In the following table the results obtained for some compounds of formula I representing the all class in comparison with erythromycin, clarithromycin and roxithromycin are reported.

Table 1

- 5 *In vitro* inhibition of IL-8 release, expressed as potency ( $IC_{50}$ ) and as efficacy (%), and of superoxide anion release ( $IC_{50}$ ).

Compound	IL-8 release		$O_2^-$ release
	potency $IC_{50} \mu M$	efficacy (%)	$IC_{50} \mu M$
1	—	29	$35.3 \pm 5.4$
2	—	29	$11.4 \pm 2.9$
5	12	$55 \pm 0.9$	>100
6	$7.7 \pm 0.3$	$37 \pm 10$	$9.9 \pm 0.69$
8	—	$19 \pm 1$	$3.5 \pm 0.2$
9	—	—	$3.5 \pm 0.5$
11	—	$22 \pm 4$	5
16	—	35	14.3
clarithromycin	$8.9 \pm 3.2$	$48 \pm 6$	$49.2 \pm 1.9$
erythromycin	—	$27 \pm 1$	>100
roxithromycin	$5.5 \pm 1.5$	$43 \pm 6$	$89 \pm 5$

#### Example 12

##### *In vivo* pharmacological activity

- 25
  - *Animals*

Male Sprague-Dawley rats weighing between 200 and 300 g were used.

The rats used in the experiments were not evidently infected. The animals were maintained in standard conditions for 7 days before being sacrificed.

- *Endotoxin administration*

-24-

LPS (*E. coli* lipopolisaccharide) (endotoxin; 055:B5 serum-type; Sigma Chemical Co., St. Louis, MO) was dissolved in saline sterile solution and was administered by intraperitoneal injection (i.p.) at a dose of 6 mg/kg (1 ml/kg). Analogously the saline solution and/or the carrier (saline solution + 0.5% Tween 20) were injected into the control animals.

5 • *Compound administration (prophylactic treatment)*

Each compound was administered by i.p. injection twice a day for 6 days, the 7<sup>th</sup> day 1 hour before and 5 hours after LPS administration. The compounds were suspended with 0.5% Tween 20 in the saline solution.

• *Bronchoalveolar lavage*

10 After 24 hours from the LPS injection rats were sacrificed with an overdose of nembutal (100 mg/kg i.p.). The trachea was incannulated and the lung was washed by instilling two 5 ml aliquots of PBS (phosphate buffer saline) at 37°C and the fluid was immediately removed. The fluid was again injected and the procedure was altogether repeated three times for each aliquot.

15 • *Cell counting and differentiation*

200 µl of BALF (bronchoalveolar lavage fluid) were diluted in 1 ml of cold water and 19 ml of Isoton. The overall number of cells was counted twice by using Contraves autolyser 800. When the total number was lower than 2000, BALFs were centrifugated at 800 rpm for 10 minutes in order to separate the cells from the supernatant. The supernatants were discharged and the cells resuspended in a small amount of PBS. For the cytological evaluation, 50 µl of the solution with the resuspended cells were centrifugated for 1 minute at 1300 rpm by using a Shandon Cytospin centrifuge. The slides were fixed in acetone and dyed with DiffQuick. The differential counts of cells were performed on each slide by counting 200 cells at random; cellular types were classified as neutrophils, eosinophils and mononuclear cells according to morphological standard criteria.

25 In the following table the results obtained with some compounds of formula I representative of the all class are reported.

Table 2

*In vivo* inhibition of neutrophilia induced by LPS after repeated administrations.


-25-

Compound	Inhibition (%)
2	-31
6	-11
9	-96
erythromycin	-46
clarithromycin	-47
roxithromycin	-50

-26-

Claims

1) A compound of formula




wherein

R is hydrogen or methyl;

15 R<sub>1</sub> and R<sub>2</sub> are both hydrogen or they together form a bond;

R<sub>3</sub> is hydrogen, a linear or branched C<sub>1</sub>-C<sub>5</sub> alkyl group, a benzyl group, optionally substituted by one or more substituents selected among nitro groups, hydroxy groups, carboxylic groups, amino groups, linear or branched C<sub>1</sub>-C<sub>5</sub> alkyl groups, C<sub>1</sub>-C<sub>4</sub> alkoxy carbonyl groups, aminocarbonyl groups or ciano groups, or a chain of formula

20



wherein

A is hydrogen or a phenyl group optionally substituted by one or two substituents selected among nitro groups, hydroxy groups, carboxylic groups, amino groups, linear or branched C<sub>1</sub>-C<sub>5</sub> alkyl groups, C<sub>1</sub>-C<sub>4</sub> alkoxy carbonyl groups, aminocarbonyl groups or ciano groups, or

25 a 5 or 6 membered heterocycle, saturated or unsaturated, containing from 1 to 3 heteroatoms selected among nitrogen, oxygen and sulphur, optionally substituted by one or two substituents selected among C<sub>1</sub>-C<sub>5</sub> alkyl groups, phenyl groups, hydroxy groups, oxo (=O) groups, nitro groups, C<sub>1</sub>-C<sub>4</sub> alkoxy carbonyl groups, aminocarbonyl groups, mono or di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl groups, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl groups;

-27-

X and Y, the same or different, are O, S, SO, SO<sub>2</sub> or NR<sub>4</sub>, in which R<sub>4</sub> is hydrogen, a linear or branched C<sub>1</sub>-C<sub>5</sub> alkyl group, a C<sub>1</sub>-C<sub>5</sub> alkoxy carbonyl group, a benzyloxy carbonyl group;

r is an integer from 1 to 6;

m is an integer from 1 to 8;

5 n is an integer from 0 to 2;

and their pharmaceutically acceptable salts;

the compounds 3'-desdimethylamino-3',4'-dehydroerythromycin A oxime and 3'-desdimethylamino-3',4'-dehydroerythromycin A 9-O-methyloxime being excluded.

2) A compound according to claim 1 in which R, R<sub>1</sub> and R<sub>2</sub> are hydrogen.

10 3) A compound according to claim 2 in which R<sub>3</sub> is a chain of formula



wherein

X, Y, A, r, m and n have the meanings reported in claim 1.

4) A compound according to claim 2 in which R<sub>3</sub> is a chain of formula

15



wherein

r is 2, m is 2 or 6, n is 1, Y is NR<sub>4</sub>, X is O or NR<sub>4</sub>, R<sub>4</sub> is hydrogen and A is phenyl or thiazolyl.

5) The compounds 3'-desdimethylamino-3',4'-dehydroerythromycin A oxime and 3'-desdimethylamino-3',4'-dehydroerythromycin A 9-O-methyloxime for use as antiinflammatories.

20 6) A pharmaceutical composition containing a pharmaceutically active amount of a compound according to claim 1 in admixture with a pharmaceutically acceptable carrier.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 July 2000 (20.07.2000)

PCT

(10) International Publication Number  
**WO 00/42055 A3**

(51) International Patent Classification<sup>7</sup>: C07H 17/08,  
A61K 31/70, A61P 29/00

(74) Agent: LONGONI, Alessandra; Zambon Group S.p.A.,  
Corp. Patent & Trademark Dept., Via Lillo Del Duca, 10,  
I-20091 Bresso (IT).

(21) International Application Number: PCT/EP00/00163

(22) International Filing Date: 12 January 2000 (12.01.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
MI99A000061 15 January 1999 (15.01.1999) IT

(71) Applicant (for all designated States except US): ZAM-  
BON GROUP S.P.A. [IT/IT]; Via della Chimica, 9,  
I-36100 Vicenza (IT).

(81) Designated States (national): AE, AL, AM, AT, AU, AZ,  
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,  
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,  
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,  
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,  
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,  
UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report

(88) Date of publication of the international search report:  
7 September 2001

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: MACROLIDES WITH ANTI-INFLAMMATORY ACTIVITY



(I)

(57) Abstract: 3'-Des-dimethylamino-9-ox-  
imino macrolides of formula (I) endowed  
with anti-inflammatory activity and their  
pharmaceutical use are described.

WO 00/42055 A3

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/00163

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07H17/08 A61K31/70 A61P29/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07H A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 254 534 A (ROBINSON WILLIAM S) 27 January 1988 (1988-01-27) cited in the application page 10, line 46 - line 51, 3'-desdimethylamino-3',4'-dehydroerythromy cin A 9-O-methyloxime, RN 118186-56-4 page 11, line 5 - line 10 claim 1 ---	1,5,6
A	US 3 928 387 A (KIERSTEAD RICHARD WIGHTMAN ET AL) 23 December 1975 (1975-12-23) cited in the application example 2, RN 53219-28-6 --- -/--	1,5,6

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

18 October 2000

Date of mailing of the international search report

03/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Held, P

## INTERNATIONAL SEARCH REPORT

Internr al Application No

PCT/EP 00/00163

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	LEMAHIEU, RONALD A. ET AL: "Glycoside cleavage reactions on erythromycin A. Preparation of erythronolide A" J. MED. CHEM. (1974), 17(9), 953-6 , XP000650897 page 954, compound 10, RN 53219-28-6 ---	1,5,6
A	EP 0 096 013 A (NUOVO CONSOR SANITAR NAZIONALE) 7 December 1983 (1983-12-07) claims 1,13,17 -----	1,5,6



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/00163

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0254534 A	27-01-1988	AU 7605587 A	28-01-1988
		DK 384387 A	25-01-1988
		FI 873128 A	25-01-1988
		HU 44439 A	28-03-1988
		PT 85405 A, B	01-08-1987
		JP 63107921 A	12-05-1988
		ZA 8705390 A	30-11-1988
-----			
US 3928387 A	23-12-1975	US 3992264 A	16-11-1976
		US 4070376 A	24-01-1978
-----			
EP 0096013 A	07-12-1983	IT 1189285 B	04-02-1988
		AT 20896 T	15-08-1986
		DE 3364696 D	28-08-1986
-----			